



Department of Health and Social Services  
Valerie J. Davidson, Commissioner

3601 C Street, Suite 540  
Anchorage, Alaska 99503 <http://dhss.alaska.gov/dph/Epi>

Division of Public Health  
Jay C. Butler, MD, Chief Medical Officer  
and Director  
Local (907) 269-8000  
24 Hour Emergency (800) 478-0084

Editors:  
Joe McLaughlin, MD, MPH  
Louisa Castrodale, DVM, MPH

Bulletin No. 16, June 23, 2016

## Chlamydia Infection Update — Alaska, 2015

### Background

*Chlamydia trachomatis* infection (CT) is the most common reportable infectious disease in the U.S. and in Alaska.<sup>1,2</sup> From 2010–2014, Alaska had the highest CT infection rate in the nation.

Often asymptomatic, untreated CT infection can cause miscarriage, pre-term labor, low birth weight,<sup>3</sup> conjunctivitis and pneumonia in neonates; pelvic inflammatory disease (PID), ectopic pregnancy, chronic pelvic pain, and infertility in women; and epididymitis and Reiter's syndrome in men. Moreover, CT can facilitate the transmission and acquisition of human immunodeficiency virus (HIV).


### Methods

Case data were obtained from the Section of Epidemiology (SOE) reportable conditions database and the Patient Reporting Investigating Surveillance Manager (PRISM). Population data were obtained from the Alaska Department of Labor and Workforce Development.

### Results

In 2015, 5,653 CT cases were reported to SOE, yielding an annual incidence rate of 766 cases per 100,000 persons, which represents a 1.5% decrease compared to 2014 (Figure 1).

**Figure 1. Chlamydia Infection Rates, by Year — Alaska and the United States, 2009–2015\***




\*The 2015 U.S. case rate is preliminary.

Of the 5,653 CT cases reported in 2015,

- 3,781 (67%) were in females;
- 4,578 (81%) were in persons aged ≤29 years;
- the highest rate occurred in persons aged 20–24 years at 3,822 cases per 100,000 persons; and
- American Indian/Alaska Natives (AI/AN), Native Hawaiian/Pacific Islanders (NH/PI), and Blacks accounted for the highest CT incidence rates, by race (Figure 2).


**Figure 2. Chlamydia Infection Rates (Cases per 100,000 Persons), by Race and Ethnicity — Alaska, 2014 and 2015\***



\*Note: 229 cases in 2014 and 147 cases in 2015 were of unknown or multiple races and are not included in this figure.

In 2015, the Northern and Southwest regions had the highest CT rates (2,151 and 1,702 cases per 100,000 persons, respectively). Compared to 2014 data, the greatest CT rate decrease occurred in the Southwest region (14% decrease, from 1,965 to 1,702 cases per 100,000 persons), while the greatest rate increase was seen in the Northern Region (4% increase, from 2,060 to 2,151 cases per 100,000 persons [Figure 3]).

**Figure 3. Chlamydia Infection Rates, by Region — Alaska, 2014 and 2015**



### Discussion

Alaska's CT infection rate decreased by 2.7% from 2013 to 2015; this coincides with the decreases in Alaska's gonorrhea and syphilis rates in 2015.<sup>4,5</sup> Maintaining an ongoing, comprehensive approach to sexually transmitted disease (STD) control is required to decrease CT rates and to reduce the associated demographic health disparities that persist in Alaska.

### Recommendations

1. Promptly treat CT-infected patients and their sex partner(s) with azithromycin 1 g PO in a single dose, OR doxycycline 100 mg PO twice daily for 7 days.<sup>6</sup>
2. Consider the use of expedited partner therapy (EPT) for sexual partners who are unable or unwilling to present for clinical evaluation.<sup>7</sup>
3. Elicit a thorough sexual history from all STD patients to include same-sex and oral/anal activities (resources in taking a complete sexual history are available at: <http://dhss.alaska.gov/dph/Epi/hivstd/Pages/history.aspx>).
4. Obtain rectal or pharyngeal specimens, as appropriate.<sup>8</sup>
5. Test all persons who are at risk for CT for other sexually transmitted diseases, including HIV and HCV.
6. Develop a partner management plan with CT-infected patients that includes timely notification of all sex partners.
7. Screen pregnant women for STDs at the first prenatal visit; repeat testing in the third trimester for those at high risk.
8. Annually screen all sexually active females aged ≤25 years and women aged >25 years with new or multiple partners.
9. Counsel patients at risk for STDs on risk-reduction strategies, including correct and consistent condom use.
10. Report CT cases and treatment to SOE within 5 working days by fax to 561-4239. Report forms are available at: <http://dhss.alaska.gov/dph/Epi/Documents/pubs/conditions/fpmSTD.pdf>

### References

1. CDC. STD Surveillance 2014. Atlanta: U.S. DHHS; 2015. Available at: <http://www.cdc.gov/std/stats14/default.htm>
2. SOE Epidemiology Bulletin. 2014 Annual (January–December) Infectious Disease Report. No. 12, June 5, 2015. Available at: <http://epibulletins.dhss.alaska.gov/Document/Display?DocumentId=36>
3. Andrews WW, et al. The Preterm Prediction Study: association of second-trimester genitourinary CT infection with subsequent spontaneous preterm birth. *Am J Obst Gynecol* 2000;183(3):662-8.
4. SOE Epidemiology Bulletin. Gonococcal Infection Update – Alaska, 2015. No. 15, May 26, 2016. Available at: [http://www.epi.alaska.gov/bulletins/docs/b2016\\_15.pdf](http://www.epi.alaska.gov/bulletins/docs/b2016_15.pdf)
5. SOE Epidemiology Bulletin. Syphilis Update – Alaska, 2015. No. 10, April 26, 2016. Available at: [http://www.epi.alaska.gov/bulletins/docs/b2016\\_10.pdf](http://www.epi.alaska.gov/bulletins/docs/b2016_10.pdf)
6. CDC. STD Treatment Guidelines, 2015. *MMWR Recomm Rep* 2015;64(No.RR-3). Available at: <http://www.cdc.gov/std/tg2015/default.htm>
7. SOE Epidemiology Bulletin. EPT Recommendations for Alaska Providers. No. 1, January 12, 2011. Available at: <http://epibulletins.dhss.alaska.gov/Document/Display?DocumentId=163>
8. SOE Epidemiology Bulletin. Extragenital Testing for Sexually Transmitted Disease. No. 19, August 26, 2015. Available at: <http://epibulletins.dhss.alaska.gov/Document/Display?DocumentId=22>